

# DAEMEN COLLEGE IMMUNIZATION FORM



SUBMISSION OPTIONS: ONLINE: [daemen.edu/healthupload](http://daemen.edu/healthupload) | FAX: 716.839.8230  
OFFICE: Wick 116, CHIP Center | MAIL: 4380 Main St. Box #104 Amherst, NY 14226

## PART 1 – STUDENT INFORMATION

<input type="checkbox"/> M <input type="checkbox"/> F <input type="checkbox"/> Other:					
LAST NAME	FIRST NAME	MIDDLE INITIAL	GENDER	DATE OF BIRTH	
<input type="checkbox"/> Undergraduate <input type="checkbox"/> Graduate <input type="checkbox"/> Transfer <input type="checkbox"/> Re-Admit			<input type="checkbox"/> SU 20 <input type="checkbox"/> FA 20 <input type="checkbox"/> SP 20	<input type="checkbox"/> Yes <input type="checkbox"/> No	
ACADEMIC LEVEL OF ENTRY (CHECK ALL THAT APPLY)			SEMESTER OF ENTRY (INCLUDE YEAR)	STUDENT-ATHLETE	
PREFERRED PHONE (INCLUDE AREA CODE)	STREET ADDRESS		CITY	STATE	ZIP CODE

## PART 2 – IMMUNIZATIONS REQUIRED NEW YORK STATE

This section must be completed and signed or stamped by health care provider or attach immunization records from previous school, health care provider or government agency. NYS PUBLIC HEALTH LAW 2165 requires college students to show proof of immunity to Measles, Mumps and Rubella. Persons born prior to 1/1/57 are exempt from this requirement unless required by their academic major to meet clinical placement requirements. If you are exempt, you must provide proof of age. NYS PUBLIC HEALTH LAW 2167 requires colleges to distribute information about meningococcal disease and vaccination to all students. \*\*Recommended vaccinations, requirements for student athletes and students participating in external learning experiences can be found below.

### MEASLES, MUMPS, RUBELLA (MMRs)

Option 1: 2 MMR Vaccinations (1st dose after 1st birthday; \*2nd dose at least 28 days later; month, day, year must be recorded)

MMR 1: \_\_\_ / \_\_\_ / \_\_\_                      \*MMR 2: \_\_\_ / \_\_\_ / \_\_\_

Option 2: List Individual Vaccination Dates (1st dose after 1st birthday; \*2nd dose at least 28 days later)

Measles 1: \_\_\_ / \_\_\_ / \_\_\_                      Measles 2\*: \_\_\_ / \_\_\_ / \_\_\_

Mumps: \_\_\_ / \_\_\_ / \_\_\_                      Rubella: \_\_\_ / \_\_\_ / \_\_\_

Option 3: List Titer Dates and Attach Lab Report (test date, result, clinical indications)

Positive Measles Titer: \_\_\_ / \_\_\_ / \_\_\_

Positive Mumps Titer: \_\_\_ / \_\_\_ / \_\_\_

Positive Rubella Titer: \_\_\_ / \_\_\_ / \_\_\_

### MENINGITIS

Option #1: Submit completed Meningococcal immunization series within the past 5 years (month, day, year must be recorded)

MCV4/Menactra (most recent): \_\_\_ / \_\_\_ / \_\_\_

Bexero (2 doses required): \_\_\_ / \_\_\_ / \_\_\_      \_\_\_ / \_\_\_ / \_\_\_

Trumemba (3 doses required): \_\_\_ / \_\_\_ / \_\_\_      \_\_\_ / \_\_\_ / \_\_\_      \_\_\_ / \_\_\_ / \_\_\_

Option #2: Review the Meningitis Fact Sheet (back) and complete the waiver form below

By signing below, I acknowledge that I have read the meningitis disease fact sheet and I am fully aware of the risks associated with the disease, availability and effectiveness of the vaccine. I have decided that I (or my child) will NOT obtain the immunization against meningococcal meningitis disease.

\_\_\_\_\_/\_\_\_\_\_/\_\_\_\_\_  
Student Signature                      Parent/Guardian Signature (if student is under 18 years of age)

**\*\*Recommended vaccinations:** Hepatitis B, Hepatitis A, Varicella, PPD, Human Papilloma Virus, Tetanus, Diptheria and Pertussis, Flu Shot.  
**Student Athletes:** required to submit physical within 6 months of first date of participation, noting clearance to participate in physical activity.  
**Clinical Students:** required to have additional vaccinations to participate (Health Services will be in touch with students when that time comes).

Provider Name (print or stamp)	Provider Signature	Provider Address & Phone Number	Date
--------------------------------	--------------------	---------------------------------	------

# Meningococcal Disease Fact Sheet

## What is meningococcal disease?

Meningococcal disease is caused by bacteria called *Neisseria meningitidis*. It can lead to serious blood infections. When the linings of the brain and spinal cord become inflamed, it is called meningitis. The disease strikes quickly and can have serious complications, including death.

Anyone can get meningococcal disease. Some people are at higher risk. This disease occurs more often in people who are:

- Teenagers or young adults
- Infants younger than one year of age
- Living in crowded settings, such as college dormitories or military barracks
- Traveling to areas outside of the United States, such as the "meningitis belt" in Africa
- Living with a damaged spleen or no spleen
- Being treated with Soliris® or, who have complement component deficiency (an inherited immune disorder)
  - Exposed during an outbreak
- Working with meningococcal bacteria in a laboratory

## What are the symptoms?

Symptoms appear suddenly – usually 3 to 4 days after a person is infected. It can take up to 10 days to develop symptoms. Symptoms may include:

- A sudden high fever
- Headache
- Stiff neck (meningitis)
- Nausea and vomiting
- Red-purple skin rash
- Weakness and feeling very ill
- Eyes sensitive to light

## How is meningococcal disease spread?

It spreads from person-to-person by coughing or coming into close or lengthy contact with someone who is sick or who carries the bacteria. Contact includes kissing, sharing drinks, or living together. Up to one in 10 people carry meningococcal bacteria in their nose or throat without getting sick.

## Is there treatment?

Early diagnosis of meningococcal disease is very important. If it is caught early, meningococcal disease can be treated with antibiotics. But, sometimes the infection has caused too much damage for antibiotics to prevent death or serious long-term problems. Most people need to be cared for in a hospital due to serious, life-threatening infections.

## What are the complications?

Ten to 15 percent of those who get meningococcal disease die. Among survivors, as many as one in five will have permanent disabilities. Complications include:

- Hearing loss
- Brain damage
- Kidney damage
- Limb amputations

## What should I do if I or someone I love is exposed?

If you are in close contact with a person with meningococcal disease, talk with your health care provider about the risk to you and your family. They can prescribe an antibiotic to prevent the disease.

## What is the best way to prevent meningococcal disease?

The single best way to prevent this disease is to be vaccinated. Vaccines are available for people 6 weeks of age and older.

Various vaccines offer protection against the five major strains of bacteria that cause meningococcal disease:

- All teenagers should receive two doses of vaccine against strains A, C, W and Y. The first dose is given at 11 to 12 years of age, and the second dose (booster) at age 16.
- It is very important that teens receive the booster dose at age 16 in order to protect them through the years when they are at greatest risk of meningococcal disease.
- Talk to your health care provider today if your teen has not received two doses of vaccine against meningococcal strains A, C, W and Y.
- Teens and young adults can also be vaccinated against the "B" strain. Talk to your health care provider about whether they recommend vaccine against the "B" strain.

Others who should receive the vaccine include:

- Infants, children and adults with certain medical conditions
- People exposed during an outbreak
- Travelers to the "meningitis belt" of sub-Saharan Africa
- Military recruits

Please speak with your health care provider if you may be at increased risk.

## What are the meningococcal vaccine requirements for school attendance?

As of September 1, 2016, children entering grades 7 and 12 must be immunized against meningococcal disease strains A, C, W and Y according to the recommendations listed above.

## Is there an increased risk for meningococcal disease if I travel?

- Meningococcal disease and outbreaks occur in the United States and around the world. The disease is more common in the "meningitis belt" of sub-Saharan Africa. The risk is highest in people who visit these countries and who have prolonged contact with local populations during an epidemic.
- To reduce your risk of illness, wash your hands often, maintain healthy habits such as getting plenty of rest and try not to come into contact with people who are sick.

---

### Learn more about meningococcal disease:

[www.cdc.gov/meningococcal/](http://www.cdc.gov/meningococcal/)

### For more information about vaccine-preventable diseases:

[www.health.ny.gov/prevention/immunization/](http://www.health.ny.gov/prevention/immunization/)



Department  
of Health